



# Higher Algebra I — Fall 2005

Gerald Hoehn

## Problem sheet 9      October 27, 2005

**Problem 1:** Let  $R$  be a commutative ring with 1. The *ring of formal power series*  $R[[x]]$  in the indeterminate  $x$  with coefficients from  $R$ , consists of formal infinite sums

$$\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \cdots.$$

The ring structure is given by extending the ring structure of the polynomial ring  $R[x]$  to “polynomials of infinite degree” in the obvious way.

(a) Prove that  $\sum_{n=0}^{\infty} a_n x^n$  is a unit in  $R[[x]]$  if and only if  $a_0$  is a unit in  $R$ .

(b) Prove that with  $R$  also  $R[[x]]$  is an integral domain.

**Problem 2:** Let  $R$  be a commutative ring with 1. Prove that a polynomial  $f \in R[x]$  is a zero-divisor in  $R[x]$  if and only if there exists a  $b \in R$  such that  $bf = 0$ .

**Problem 3:** Let  $D$  be a squarefree integer (i.e. an integer which has no integer square besides 1 as a divisor). Let

$$\omega = \begin{cases} \sqrt{D}, & \text{if } D \equiv 2, 3 \pmod{4}, \\ \frac{1+\sqrt{D}}{2}, & \text{if } D \equiv 1 \pmod{4}. \end{cases}$$

The set  $\mathcal{O} = \{a + b\omega \mid a, b \in \mathbf{Z}\}$  is a subring of the real numbers (if  $D$  is positive) or complex numbers (if  $D$  is negative). Define the field norm of  $x + y\sqrt{D}$ , where  $x, y \in \mathbf{Q}$  (or  $\mathbf{R}$ ), by  $N(x + y\sqrt{D}) = (x + y\sqrt{D})(x - y\sqrt{D}) = x^2 - Dy^2$ .

Prove that for  $D = -2, -3, -7$  or  $-11$  the ring  $\mathcal{O}$  is an Euclidean Domain with respect to  $N$ . (Hint: Modify the proof for  $\mathbf{Z}[i]$  ( $D = -1$ ) given in the lecture. For  $D = -3, -7, -11$ , prove that every element of  $\mathbf{C}$  differs from an element of  $\mathcal{O}$  by an element whose norm is at most  $(1 + |D|)^2 / (16|D|)$ , which is less than 1 for these values of  $D$ . Plotting the points of  $\mathcal{O}$  in  $\mathbf{C}$  may be helpful.)

**Problem 4:** Let  $K$  be a field, let  $a_1, \dots, a_n$  be pairwise different and let  $b_1, \dots, b_n$  be arbitrary elements of  $K$ . Show that there is a unique polynomial  $f \in K[x]$  with  $\deg(f) \leq n - 1$  and  $f(a_i) = b_i$  for  $i = 1, \dots, n$ .